

Topological Excitations in Double-Layer Quantum Hall systems

Kyungsun Moon

Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019

(February 1, 2008)

Abstract

Double-layer quantum Hall systems with spontaneous broken symmetry can exhibit a novel manybody quantum Hall effect due to the strong interlayer coherence. When the layer separation becomes close to the critical value, quantum fluctuations can destroy the interlayer coherence and the quantum Hall effect will disappear. We calculate the renormalized isospin stiffness ρ_s due to quantum fluctuations within the Hartree-Fock-RPA formalism. The activation energy of the topological excitations thus obtained demonstrates a nice qualitative agreement with recent experiment.

PACS numbers: 75.10.-b, 73.20.Dx, 64.60.Cn

Double-layer quantum Hall systems (DLQHS) have drawn much theoretical and experimental attention recently [1–8]. When the distance d between the layers is comparable to the mean particle spacing, strong interlayer correlations induce a novel manybody quantum Hall effect at $\nu_T = 1/m$ (m is an odd integer), where ν_T is the total filling factor [3,8,10]. Since each layer at $\nu = 1/(2m)$ alone does not support quantum Hall effect, it manifests a *genuine* manybody correlation effect. When the layer separation becomes large and close to the critical value, quantum fluctuations can destroy the interlayer coherence and accordingly quantum Hall effect will disappear [3,10,9]. By mapping the layer degrees of freedom onto the $S = \frac{1}{2}$ isospin variables, the interlayer correlations can be quantified as isospin stiffness ρ_s . Since most of the recent experimental systems [3,10] belong to the quantum fluctuation dominated regime, it is *crucial* to obtain the reliable estimate of the renormalized isospin stiffness ρ_s which incorporates the effect of quantum fluctuations.

In this Letter, we calculate the renormalized isospin stiffness ρ_s and the effective tunneling amplitude t_R within the Hartree-Fock-RPA formalism. By taking into account the important vertex corrections [7,9,12], we explicitly show that ρ_s vanishes at $d/\ell \cong 1$, where $\ell = \sqrt{\hbar c/|e|B}$ is the magnetic length. The activation energy of the lowest charge excitations (*e.g.* Meron pairs *or* domain-wall string solitons) is estimated and shows a nice qualitative agreement with the recent experiment [10]. We consider the DLQHS with finite tunneling t and parallel magnetic field $B_{\parallel}\hat{y}$. We concentrate on the $\nu_T = 1$ quantum Hall effect and assume that the electrons are lying on the lowest Landau level due to the strong perpendicular magnetic field $B_{\perp}\hat{z}$ and the real spins are fully spin-polarized due to the finite Zeeman gap. With the choice of the following gauge $\mathbf{A} = (0, B_{\perp}x, -B_{\parallel}x)$, the Hamiltonian of the system can be written by

$$\begin{aligned} \mathcal{H} = & -te^{-Q^2/4} \sum_X \left(e^{iQX} c_{X\uparrow}^{\dagger} c_{X\downarrow} + e^{-iQX} c_{X\downarrow}^{\dagger} c_{X\uparrow} \right) \\ & + \frac{1}{2} \sum_{\sigma, \sigma'} \sum_{X_1, X_2, X' \neq 0} V_{\sigma\sigma'}(X', X_1 - X_2) c_{X_1 - X'\sigma}^{\dagger} c_{X_2\sigma'}^{\dagger} c_{X_2 - X'\sigma'} c_{X_1\sigma} \end{aligned} \quad (1)$$

where $\sigma = +1(-1)$ represents an electron in the upper(lower) layer, X the guiding center coordinate, and $Q = dB_{\parallel}/\ell^2 B_{\perp}$. With the following convention of the units $e = \ell = \hbar = 1$,

$V_{\sigma\sigma'}$ is an intralayer interaction V_A for $\sigma = \sigma'$ and an interlayer interaction V_E for $\sigma \neq \sigma'$: $\tilde{V}_A = 2\pi/\epsilon q$, $\tilde{V}_E = 2\pi e^{-qd}/\epsilon q$, and

$$V_{\sigma\sigma'}(X, X_1 - X_2) = \frac{1}{L^2} \sum_{q_y} e^{-q^2/2} \tilde{V}_{\sigma\sigma'}(X, q_y) e^{iq_y(X_1 - X_2)} \quad (2)$$

where $q = \sqrt{X^2 + q_y^2}$. We notice that $B_{\parallel}$ induces an Aharonov-Bohm phase $e^{\pm iQX}$ depending on the sense of interlayer tunneling.

The relatively weak interlayer interaction comparing to the intralayer interaction suppresses local density fluctuations. Since this corresponds to the m_z^2 -term in the magnetic analogy, the isospins are forced to lie on the $\hat{x} - \hat{y}$ isospin space [6]. Hence the DLQHS can be viewed as quantum XY -ferromagnets. We obtain the following energy functional

$$E[\theta] = \int d^2r \left\{ \frac{1}{2} \rho_s (\nabla \theta)^2 - \frac{t}{2\pi} \cos [\theta(\mathbf{r}) - Qx] \right\} \quad (3)$$

where $\theta(\mathbf{r})$ represents the isospin orientation. This is precisely the Pokrovsky-Talapov (PT) model [15], which exhibits a commensurate-incommensurate transition. For small Q and hence $B_{\parallel}$, the phase obeys $\theta(\mathbf{r}) = Qx$. As $B_{\parallel}$ increases, the local field tumbles too rapidly and a continuous phase transition to an incommensurate state with broken translation symmetry occurs [2,6]. The ground state energy of the commensurate state is given by

$$\epsilon(Q, t) \equiv \frac{E[Q, t]}{A} = \frac{1}{2} \rho_s Q^2 - \frac{t}{2\pi} e^{-Q^2/4} \quad (4)$$

where ρ_s measures the interlayer coherence. By imposing the following condition $\sqrt{t/\rho_s} \gg Q > 0$, one can guarantee that the ground state of the system is the commensurate one and ρ_s can be obtained by the following procedure

$$\rho_s = \lim_{t \rightarrow 0} \lim_{Q \rightarrow 0} \frac{d^2 \epsilon(Q, t)}{dQ^2}. \quad (5)$$

An appealing feature of the commensurate state is that the self-consistent Hartree-Fock ground state is also an eigenstate of the tunneling Hamiltonian alone. One can easily diagonalize the tunneling Hamiltonian in terms of the α_X, β_X -operators

$$H_t = -te^{-Q^2/4} \sum_X (\alpha_X^\dagger \alpha_X - \beta_X^\dagger \beta_X) \quad (6)$$

where α_X, β_X correspond to the symmetric and antisymmetric state with respect to the spatially varying tumbling field: $\alpha_X = (e^{-iQX/2}c_{X\uparrow} + e^{iQX/2}c_{X\downarrow})/\sqrt{2}$, $\beta_X = (e^{-iQX/2}c_{X\uparrow} - e^{iQX/2}c_{X\downarrow})/\sqrt{2}$. The ground state $|\Phi\rangle$ is a completely filled state of the α_X -particles. Since $|\Phi\rangle$ is an eigenstate of the full Hartree-Fock Hamiltonian *too*, it is straightforward to calculate the Hartree-Fock ground state energy yielding

$$\begin{aligned}\frac{E_{HF}}{A} &= \frac{1}{A} \langle \Phi | \mathcal{H} | \Phi \rangle \\ &= -\frac{t}{2\pi} e^{-Q^2/4} - \frac{1}{4\pi} \int \frac{dX dq_y}{(2\pi)^2} e^{-q^2/2} \tilde{V}_o^Q(X, q_y)\end{aligned}\quad (7)$$

where $\tilde{V}_o^Q(X, q_y) = (\tilde{V}_A + \tilde{V}_E \cos QX)/2$. Using Eq.(5), we obtain the Hartree-Fock estimate of isospin stiffness

$$\rho_s^0 = \frac{1}{32\pi^2} \int_0^\infty dq \tilde{V}_E(q) e^{-q^2/2} q^3. \quad (8)$$

Now we want to calculate the fluctuation corrections to ρ_s^0 using the Hartree-Fock-RPA scheme. The Green function $G_{XX'}^{\sigma\sigma'}(\tau - \tau')$ is defined as follows

$$G_{XX'}^{\sigma\sigma'}(\tau - \tau') \equiv \langle T_\tau C_{\sigma X}(\tau) C_{\sigma' X'}^\dagger(\tau') \rangle. \quad (9)$$

The retarded Green function $G_{XX'}^{\sigma\sigma'}(\omega)$ of the Hartree-Fock mean field Hamiltonian is given by

$$G_{XX'}^{\sigma\sigma'}(\omega) = \delta_{X,X'} \frac{1}{2} e^{\frac{i}{2}(\sigma - \sigma')QX} \left\{ \frac{1}{\omega - \epsilon_\alpha - i\eta} + \sigma\sigma' \frac{1}{\omega - \epsilon_\beta + i\eta} \right\} \quad (10)$$

where $\epsilon_\alpha = -te^{-Q^2/4} - \sum_X \langle 0X | \tilde{V}_0^Q | X0 \rangle$, $\epsilon_\beta = te^{-Q^2/4} - \sum_X \langle 0X | \tilde{V}_x^Q | X0 \rangle$, and $\tilde{V}_x^Q(X, q_y) = (\tilde{V}_A - \tilde{V}_E \cos QX)/2$. $\delta_{X,X'}$ indicates that the system is translationally invariant along the $\hat{y}$ -direction in our chosen gauge and we only keep a single index X from now on. The RPA ground state energy can be calculated as follows [14]

$$E_{\text{RPA}} = \sum_{n=2}^{\infty} \frac{i}{2n} \langle \Phi | (D V)^n | \Phi \rangle \quad (11)$$

where D is the Hartree-Fock particle-hole propagator defined by

$$\begin{aligned}
D_{XX'}^{\sigma\sigma'}(\omega) &= \int \frac{d\omega'}{2\pi} G_X^{\sigma\sigma'}(\omega + \omega') G_{X'}^{\sigma'\sigma}(\omega') \\
&= \frac{i}{4} \sigma\sigma' e^{\frac{i}{2}(\sigma-\sigma')Q(X-X')} A(\omega)
\end{aligned} \tag{12}$$

where $\Delta_x(Q) = \Delta_{\text{SAS}} e^{-Q^2/4} + \sum_X \langle 0X | \tilde{V}_E \cos QX | X0 \rangle$ represents the Q -dependent exchange-enhanced gap with $\Delta_{\text{SAS}} = 2t$ and $A(\omega) = (\frac{1}{\omega - \Delta_x(Q) + i\eta} - \frac{1}{\omega + \Delta_x(Q) - i\eta})$. The strong enhancement of the tunneling gap due to the Coulomb exchange energy stabilizes our perturbative analysis. When σ and σ' are opposite to each other, $D_{XX'}^{\sigma\sigma'}$ represents an interlayer tunneling process which picks up an Aharonov-Bohm phase of $e^{\pm iQ(X-X')}$. We first consider the n -th order bubble diagram (refer to Fig.(1))

$$\begin{aligned}
E_{\text{RPA}}^{(n)} &= \frac{(-i)^{n-1}}{2n} \sum_{\{\sigma_i, X_i\}} \int \frac{d\omega}{2\pi} D_{X_1 X_2}^{\sigma_{2n} \sigma_1}(\omega) \cdots D_{X_{2n-1} X_{2n}}^{\sigma_{2n-2} \sigma_{2n-1}}(\omega) \\
&\quad \langle X_1 X_4 | V_{\sigma_1 \sigma_2} | X_2 X_3 \rangle \cdots \langle X_{2n-1} X_2 | V_{\sigma_{2n-1} \sigma_{2n}} | X_{2n} X_1 \rangle.
\end{aligned} \tag{13}$$

We notice that the momentum transfer $X = X_{2i-1} - X_{2i}$ is conserved. The sum over isospins can be performed using the following trick. Since X stays the same for all the bubbles, the phase factor in $D_{X_1 X_2}^{\sigma_{2n} \sigma_1}$ can be rearranged as follows

$$D_{X_1 X_2}^{\sigma_{2n} \sigma_1}(\omega) \cdots D_{X_{2n-1} X_{2n}}^{\sigma_{2n-2} \sigma_{2n-1}}(\omega) = [iA(\omega)]^n \prod_{i=1}^n \left[\frac{\sigma_{2i-1} \sigma_{2i}}{4} e^{-\frac{i}{2}(\sigma_{2i-1} - \sigma_{2i})QX} \right]. \tag{14}$$

Using Eq.(14), the isospin-sum can be factorized and we obtain the following useful relations

$$\frac{1}{4} \sum_{\sigma_1, \sigma_2} \sigma_1 \sigma_2 e^{-\frac{i}{2}(\sigma_1 - \sigma_2)QX} \langle X_1 X_4 | V_{\sigma_1 \sigma_2} | X_2 X_3 \rangle = V_x^Q(X, X_1 - X_3). \tag{15}$$

Since our gauge choice conserves the momentum along the $\hat{y}$ -direction, we finally obtain

$$\frac{E_{\text{RPA}}^{(n)}}{A} = \frac{i}{2n} \frac{1}{L^2} \sum_{X, q_y} \int \frac{d\omega}{2\pi} \left[\frac{1}{2\pi} e^{-q^2/2} \tilde{V}_x^Q(X, q_y) A(\omega) \right]^n. \tag{16}$$

The sum of $E_{\text{RPA}}^{(n)}$ with respect to n yields

$$\frac{E_{\text{RPA}}}{A} = \frac{i}{2} \frac{1}{L^2} \sum_{X, q_y} \int \frac{d\omega}{2\pi} \left\{ -\ln \left[1 - \frac{1}{2\pi} e^{-q^2/2} \tilde{V}_x^Q(X, q_y) A(\omega) \right] - \frac{1}{2\pi} e^{-q^2/2} \tilde{V}_x^Q(X, q_y) A(\omega) \right\}. \tag{17}$$

The integration over frequency ω can be explicitly performed and E_{RPA} can be written by

$$\begin{aligned} \frac{E_{\text{RPA}}}{A} = & \frac{1}{2L^2} \sum_{X,q_y} \left\{ \sqrt{\Delta_x^2(Q) + \frac{1}{\pi} e^{-q^2/2} \tilde{V}_x^Q(X, q_y) \Delta_x(Q)} - \Delta_x(Q) \right\} \\ & - \frac{1}{2L^2} \sum_{X,q_y} \frac{1}{2\pi} e^{-q^2/2} \tilde{V}_x^Q(X, q_y). \end{aligned} \quad (18)$$

The last term in Eq.(18) plus E_{HF} does not have a Q -dependence. By expanding both $\Delta_x(Q)$ and $\tilde{V}_x^Q$ upto the quadratic order in Q , we obtain

$$\begin{aligned} \rho_s = & \int \frac{d^2q}{(2\pi)^2} \left\{ \frac{\frac{1}{16\pi} e^{-q^2/2} q^2 \tilde{V}_E \Delta_{x0} - 2e^{-q^2/2} \rho_s^0 \tilde{V}_x}{\sqrt{\Delta_{x0}^2 + \frac{1}{\pi} e^{-q^2/2} \tilde{V}_x \Delta_{x0}}} \right. \\ & \left. + \frac{4\pi \rho_s^0 \left(\sqrt{\Delta_{x0}^2 + \frac{1}{\pi} e^{-q^2/2} \tilde{V}_x \Delta_{x0}} - \Delta_{x0} \right)}{\sqrt{\Delta_{x0}^2 + \frac{1}{\pi} e^{-q^2/2} \tilde{V}_x \Delta_{x0}}} \right\} \end{aligned} \quad (19)$$

where $\tilde{V}_x = (\tilde{V}_A - \tilde{V}_E)/2$ and $\Delta_{x0} = \Delta_x(Q=0)$.

It is well-known that the coherent double-layer quantum Hall state is unstable due to the presence of a charge density wave(CDW) instability at a finite wave-vector $k\ell \cong 1$ for $d/\ell \geq 1.2$ [7,9,12]. In order to correctly capture this important quantum fluctuations, we take into account the vertex corrections, *i.e.*, a sum of ladders in the polarization insertions in Fig.(1). It amounts to replacing $\Delta_x(Q)$ with the correct dispersion relation $\omega(\mathbf{q}, Q)$ of the collective mode in Eq.(18)

$$\omega(\mathbf{q}, Q) = \sqrt{D_z(\mathbf{q}, Q) D_y(\mathbf{q}, Q)} \quad (20)$$

where $D_z(\mathbf{q}, Q)$ and $D_y(\mathbf{q}, Q)$ are given by

$$D_z(\mathbf{q}, Q) = \Delta_{\text{SAS}} + \frac{1}{\pi} \tilde{V}_x^Q e^{-q^2/2} + \int \frac{d^2k}{(2\pi)^2} \tilde{V}_E(k) \cos Qk_y e^{-k^2/2} - \int \frac{d^2k}{(2\pi)^2} \tilde{V}_A(k) e^{-k^2/2} e^{i\mathbf{k} \times \mathbf{q} \cdot \hat{z}} \quad (21)$$

$$D_y(\mathbf{q}, Q) = \Delta_{\text{SAS}} + \int \frac{d^2k}{(2\pi)^2} \tilde{V}_E(k) \cos Qk_y e^{-k^2/2} - \int \frac{d^2k}{(2\pi)^2} \tilde{V}_E(k) \cos Qk_y e^{-k^2/2} e^{i\mathbf{k} \times \mathbf{q} \cdot \hat{z}}. \quad (22)$$

Fig.(2) demonstrates that ρ_s obtained by RPA with vertex corrections indeed exhibits a dramatic reduction from the Hartree-Fock *or* RPA estimates. One can also notice that ρ_s vanishes at $d/\ell \cong 1$ due to the proximity to the strong CDW instability. Quantum

fluctuations can also reduce the normalized magnetization $\langle m_x \rangle$ —the *order parameter* of the DLQHS. One can calculate $\langle m_x \rangle$ using the following procedure [6]

$$\begin{aligned} \langle m_x \rangle &= -\lim_{t \rightarrow 0} \lim_{Q \rightarrow 0} 2\pi \frac{d\epsilon(Q, t)}{dt} \\ &= 1 - \int \frac{d^2 q}{2\pi} \left\{ \frac{\Delta_{x0} + \frac{1}{2\pi} e^{-q^2/2} \tilde{V}_x}{\sqrt{\Delta_{x0}^2 + \frac{1}{\pi} e^{-q^2/2} \tilde{V}_x \Delta_{x0}}} - 1 \right\}. \end{aligned} \quad (23)$$

The renormalization of $\langle m_x \rangle$ implies that the effective tunneling amplitude t_R is reduced by a factor of $\langle m_x \rangle$: $t_R = t \langle m_x \rangle$. The inset of Fig.(2) shows that $\langle m_x \rangle$ via RPA with vertex corrections is strongly reduced from the Hartree-Fock *or* RPA estimates. Based on the general field-theoretical point of view, we speculate that in the absence of tunneling, $\langle m_x \rangle$ will vanish at the same value of d as ρ_s [6]. Here $\langle m_x \rangle$ vanishes at $d/\ell \sim 0.7$, which is comparable to $d/\ell \sim 1$ from ρ_s .

The activation energy of the lowest charge excitations can be calculated from ρ_s and t_R obtained above. The subscript R is omitted from now on. When the tunneling is very weak, *i.e.*, $\tilde{t} \leq 2.4 \times 10^3 \tilde{\rho}_s^3$, the lowest charge excitations are meron-pairs, where $\tilde{t}, \tilde{\rho}_s$ are measured in units of $e^2/\epsilon\ell$ [6,13]. Since the merons carry charge $\pm \frac{1}{2}e$ depending on the vorticity and the core-spin configurations, the energy E_{MP} of meron pairs with charge $\pm e$ can be determined by balancing the Coulomb repulsion and the logarithmic attraction: $E_{MP} \sim 4\pi\rho_s$ [6]. As t increases *or* ρ_s decreases, it becomes too costly to flip the isospins over a broad area due to the large tunneling energy cost. Hence the meron-pairs are confined by a linear string tension $T_0 = 8(t\rho_s/2\pi)^{1/2}$ and the lowest charge excitations become domain-wall string solitons(DWS) [6,16]. The activation energy of the DWS can be estimated by balancing the Coulomb repulsion and linear string tension: $E_{DWS} \sim (e^2 T_0/\epsilon)^{1/2}$ [6,16]. We have chosen $t/(e^2/\epsilon\ell) \cong 4 \times 10^{-3}$ [17]. Fig.(3) shows the d/ℓ -dependence of the activation energy. Since one needs to create a charge *neutral* excitation at $\nu_T = 1$, E_A is multiplied by a factor of two. Below $d/\ell \sim 0.7$, the meron pairs are the lowest charge excitations and above are the DWS. The theoretical result thus obtained demonstrates a nice qualitative agreement with the experiment [10]. When d/ℓ vanishes, one recovers a skyrmion-antiskyrmion pair energy $8\pi\rho_s$ [18]. In real experimental systems, when the layer separation becomes small,

the interlayer tunneling becomes much easier and the system acts like a single wide quantum well. As the Zeeman gap is much smaller than Δ_{SAS} , the real spin degrees of freedom will become important and the lowest charge excitations are *real spin*-textured quasiparticles.

To summarize, we have calculated the renormalized isospin stiffness ρ_s and the effective tunneling amplitude t_R within the Hartree-Fock-RPA formalism. We have explicitly shown that ρ_s vanishes at $d/\ell \cong 1$. We have also estimated the d/ℓ -dependent activation energy of the topological excitations.

ACKNOWLEDGMENTS

It is my pleasure to acknowledge useful conversations with S.M. Girvin, A.H. MacDonald, K. Mullen, and S.Q. Murphy. The work was supported by NSF DMR-9502555. I want to acknowledge the Aspen center for physics where part of this work has been performed.

REFERENCES

- [1] S.M. Girvin and A.H. MacDonald, to appear in *Novel Quantum Liquids in Low-Dimensional Semiconductor Structures*, edited by S. Das Sarma and A. Pinczuk (Wiley, New York, 1996).
- [2] Kun Yang, K. Moon, L. Zheng, A.H. MacDonald, S.M. Girvin, D. Yoshioka, and Shou-Cheng Zhang, *Phys. Rev. Lett.* **72**, 732 (1994).
- [3] S.Q. Murphy, J.P. Eisenstein, G.S. Boebinger, L.N. Pfeiffer, and K.W. West, *Phys. Rev. Lett.* **72**, 728 (1994).
- [4] X.G. Wen and A. Zee, *Phys. Rev. Lett.* **69**, 1811 (1992).
- [5] Z.F. Ezawa and A. Iwazaki, *Int. J. of Mod. Phys. B* **19**, 3205 (1992).
- [6] K. Moon *et al.*, *Phys. Rev. B* **51**, 5138 (1995); Kun Yang *et al.*, *Phys. Rev. B* **54**, 11644 (1996).
- [7] H.A. Fertig, *Phys. Rev. B* **40**, 1087 (1989).
- [8] T.S. Lay *et al.*, *Phys. Rev. B* **50**, 17725 (1994).
- [9] A.H. MacDonald, P.M. Platzman, and G.S. Boebinger, *Phys. Rev. Lett.* **65**, 775 (1990).
- [10] R.J. Hyndman *et al.*, *preprint* (Univ. of Nottingham).
- [11] Tin-Lun Ho, *Phys. Rev. Lett.* **73**, 874 (1994).
- [12] R. Côté, L. Brey, H.A. Fertig, and A.H. MacDonald, *Phys. Rev. B* **51**, 13475 (1995); R. Côté and A.H. MacDonald, *Phys. Rev. B* **44**, 8759 (1991).
- [13] K. Yang and A.H. MacDonald, *Phys. Rev. B* **51**, 17247 (1995).
- [14] J. W. Negele and H. Orland, *Quantum Many-Particle Systems* (Addison-Wesley 1987).
- [15] Per Bak, *Rep. Prog. Phys.* **45**, 587 (1982).


- [16] N. Read, *Phys. Rev. B* **52**, 1926 (1995).
- [17] We notice that $\langle m_x \rangle$ has *much* stronger dependence on the weak tunneling than ρ_s .
- [18] S.L. Sondhi, A. Karlhede, S.A. Kivelson, and E.H. Rezayi, *Phys. Rev. B* **47**, 16419 (1993).

FIGURES


FIG. 1. The n -th order RPA bubble diagram.

FIG. 2. The renormalized isospin stiffness ρ_s : the solid line stands for the Hartree-Fock result ρ_s^0 , the dashed line for the RPA, and the long-dashed line for the RPA with the vertex corrections. The inset shows the corresponding renormalized magnetizations $\langle m_x \rangle$.


FIG. 3. The activation energy of the lowest charge excitations.



K. Moon Fig.3



K. Moon Fig. 2



K. Moon Fig.1